

Evidence for Infections by the Same Strain of Beta 2-toxigenic *Clostridium perfringens* Type A Acquired in One Hospital Ward

DOMINIKA SALAMON¹, DOROTA OCHOŃSKA¹, ILONA WOJAK², EWA MIKOŁAJCZYK²,
MAŁGORZATA BULANDA³ and MONIKA BRZYCHCZY-WŁOCH¹

¹ Department of Molecular Medical Microbiology, Chair of Microbiology, Faculty of Medicine,
Jagiellonian University Medical College, Krakow, Poland

² Department of Microbiological Diagnostics, Blessed Father Jerzy Popieluszko Provincial Specialist Hospital,
Włocławek, Poland

³ Department of Epidemiology of Infections, Chair of Microbiology, Faculty of Medicine,
Jagiellonian University Medical College, Krakow, Poland

Submitted 27 February 2019, revised 30 April 2019, accepted 21 May 2019

Abstract

This study conducts a comparative phenotypic and genetic analysis of *C. perfringens* strains isolated from two patients hospitalized at the same time in 2017 in the surgical ward of the Provincial Specialist Hospital in Włocławek (Kujawsko-Pomorskie Province) who developed necrotizing soft tissue infections (NSTI). To explain the recurring cases of this infection, a comparative analysis was performed for these strains and the ones originating from infections recorded at the same hospital in three patients with gas gangrene in 2015. The two *C. perfringens* isolates studied in 2017 (8554/M/17 from patient No. 1 and 8567/M/17 from patient No. 2) had identical biochemical profiles. A comparison of research results using multiplex PCR from 2017 with a genetic analysis of strains from 2015 enabled us to demonstrate that the strains currently studied have the genes encoding the same toxins (α and β_2) as the two strains analyzed in 2015: no. 7143 (patient No. 3) and no. 7149 (patient No. 2). A comparative analysis of the strain profiles obtained with pulsed-field gel electrophoresis (PFGE) in 2017 with the results from 2015 has found one identical and genetically unique restriction profile, corresponding to one clone of *C. perfringens* comprising of two strains: no. 8567/M/17 (patient No. 2 in 2017) and no. 7143 (patient No. 3 in 2015). The epidemiological data and detailed analysis of the course of both events suggest that this clone of *C. perfringens* possibly survived in adverse conditions of the external environment in the operating block of this hospital for many months.

Key words: *Clostridium perfringens*, beta 2 (β_2)-toxin, NSTI, molecular diagnostics

Introduction

Clostridium perfringens, a Gram-positive, anaerobic bacillus, is a bacterium commonly found in nature. Its presence can be confirmed in both the external environment (water, soil, and sewage) as well as in the digestive tract of humans and animals, where it is part of the microbiome. The important carriers of this microbe are the elderly and people engaged in the processing and distribution of food. Nonetheless, its ability to produce numerous toxins and enzymes as well as to form spores makes this bacterium a dangerous pathogen of humans and animals (Kędzińska et al. 2012).

The scientific literature indicates several toxins produced by *C. perfringens* that play a vital role in the pathogenicity of this bacterium, they are: alpha (α , *C. perfringens* alpha toxin – CPA) and a synergistic theta, i.e. perfringolysin (θ , perfringolysin O – PFO), as well as beta (β , *C. perfringens* beta toxin – CPB), epsilon (ϵ , epsilon toxin – ETX), iota (ι , iota toxin – ITX), enterotoxin (*C. perfringens* enterotoxin – CPE), necrotic enteritis B-like toxin (NetB), and β_2 toxin, discovered in 1997, the role of which is not fully explained and still requires further research (van Asten et al. 2010; Brzychczy-Włoch and Bulanda 2014; Navarro et al. 2018). It is known that β_2 -toxin is produced by animal strains of

* Corresponding author: M. Brzychczy-Włoch, Department of Molecular Medical Microbiology, Chair of Microbiology, Faculty of Medicine, Jagiellonian University Medical College, Krakow, Poland; e-mail: m.brzychczy-wloch@uj.edu.pl

© 2019 Dominika Salamon et al.

This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

C. perfringens, especially in the course of necrotic enteritis in pigs. Studies on human strains of this bacterial species, isolated from patients with food poisoning, an antibiotic-associated diarrhea and sporadic diarrhea, but also from healthy carriers, have shown that, in some cases, they can have *cpb2* gene for β 2-toxin (Johansson et al. 2006, van Asten et al. 2010; Allaart et al. 2014).

Owing to their ability to produce various toxins, strains belonging to the species *C. perfringens* were divided into 6 toxinotypes, A – G, which are responsible for gastrointestinal tract infections in humans (types A, C, F) and animals (types B, C, D, E, G) as well as severe soft tissue infections in humans and animals (type A) (Stevens et al. 2012; Navarro et al. 2018; Rood et al. 2018).

The term *necrotizing soft tissue infections* (NSTI) comprises a group of diseases (especially necrotizing fasciitis – NF) causing rapid and extensive soft tissue necrosis, which often leads to systemic infection, shock, and multiple organ failure, and ultimately, death (Stevens et al. 2017). These infections have different causes, risk factors, location, and pathomechanisms. NSTIs are often divided into two types. Type 1 is a multi-bacterial infection, the significant parts of which are anaerobic bacteria (among others, from the genus *Bacteroides* or *Clostridium*) and facultative anaerobes (among others of the family Enterobacteriaceae). It is often diagnosed in the elderly and the risk factors are diabetes, bedsores, hemorrhoids or anal surgery, and urological surgery or gynecological procedures. A peculiar infection of this type is a Fournier gangrene, which may develop as secondary to damaged mucous membranes of the gastrointestinal tract or urinary tract (Žaba et al. 2009; Stevens et al. 2017; Kuzaka et al. 2018). *C. perfringens* is listed as one of the numerous bacteria identified in the course of this infection. Although in recent years, its share in the infection seems to be getting smaller (Kuzaka et al. 2018), the literature still describes cases of Fournier gangrene in which *C. perfringens* was cultured among the other infectious agents (Wróblewska et al. 2014; Stevens et al. 2017). Type 2 of NSTI is usually associated with infection with a single bacterial species (e.g. MRSA) and often affects limbs. Some people distinguish Type 3, i.e. infection caused by a particular bacterial species: *Aeromonas hydrophila*, *Vibrio vulnificus* or a species from the genus *Clostridium*, which is most often isolated from gas gangrene cases (Stevens et al. 2017).

From the point of view of epidemiology and future management in cases of recurring NSTI in the same hospital, it is important to identify the differences and similarities among bacterial strains from every patient. The use of molecular methods allows for fast and precise identification of the bacteria isolated. The objective of this study is a comparative phenotypic and genetic analysis of *C. perfringens* strains isolated from two

patients hospitalized at the same time in 2017 in the Provincial Specialist Hospital in Włocławek (Kujawsko-Pomorskie Province) who developed soft tissue infections. Trying to explain recurring cases of this infection, we also carried out an additional comparative analysis of *C. perfringens* strains isolated in 2017 with the strains isolated in 2015 from three patients with gas gangrene in the same hospital that have been already described by our team (Brzychczy-Włoch et al. 2016).

Experimental

Materials and Methods

The study involved the microbiological analysis of *C. perfringens* strains isolated from biological specimens originating from two patients with NSTI who were hospitalized between 17th May 2017 and 4th July 2017 in a particular Department of General Surgery in the hospital in Włocławek. The analysis of microbiological and epidemiological data concerning three patients hospitalized in the same hospital between 15th April 2015 and 20th April 2015 (Brzychczy-Włoch et al. 2016) was also performed.

Source of the isolates and epidemiological data from 2017. Patient 1. A man, aged 60, admitted on 17.05.2017 to the General Surgery Department due to critical vascular insufficiency of the lower limbs. Due to the ischemia of the lower skin flap covering the stump and inflammatory infiltration in the postoperative wound on 31.05.2017, the stump wound was swabbed. The growth of *C. perfringens* on microbiological media under anaerobic conditions and of coagulase-negative staphylococci (CNS) under aerobic conditions was demonstrated. Control swabs from the healing wound were microbiologically negative. The microbiological examination of the blood taken from the patient did not show bacterial growth.

Patient 2. A man, aged 60, admitted urgently on 31.05.2017 to the General Surgery Department due to anorectal abscess. In the course of the diagnostics undertaken, disseminated sigmoid colon cancer was eventually diagnosed. Past medical history revealed a stay in the same unit in May 2014, when the patient underwent surgery due to right-sided incarcerated inguinal hernia.

During the relevant hospitalization, on 31.05.2017, abscess contents were collected from the patient for microbiological diagnostics. Microbiological testing of the purulent content detected the growth of *Bacteroides fragilis* and *C. perfringens* on microbiological media under anaerobic conditions, and *Escherichia coli* and *Pseudomonas aeruginosa* under aerobic conditions. Microbiological testing of the patient's blood did not show bacterial growth. During the microbiological

examination of a rectal swab carried out on 6.06.2017, growth of *C. perfringens* was detected.

The prevention of epidemic outbreak and epidemiological investigations were initiated. Contact isolation was employed with both patients. Lavasepsis was used for wound dressing with 0.9% NaCl solution and superoxidized solution in the form of Microdacyn (Oculus). A sporicidal agent, Incidin Active (Ecolab), was applied for surface disinfection in the ward. When microbiological results were obtained, the entire General Surgery Department and the Main Operating Block were covered by the control. Growth of *C. perfringens* was not demonstrated in cultures collected from the wounds of other patients, the skin of the staff's hands, or on the surfaces in all rooms and on the tools.

Microbiological diagnostics of *C. perfringens* strains. Samples of two patients were collected during a routine check-up by medical staffs and were diagnosed in the Department of Microbiological Diagnostics of the Provincial Specialist Hospital in Włocławek. Specimens from the patients (wound swab containing activated carbon (COPAN) from patient No. 1 and abscess contents from patient No. 2) were put into sterile test tubes with a transport medium. The culture was carried out on Columbia Agar (BioMérieux) as well as in the fluid thioglycollate medium with resazurin (BioMaxima). Solid media with the inoculated materials were incubated under aerobic conditions at 37°C for 48 h and under anaerobic conditions at 37°C for 24 h, and the liquid medium under aerobic conditions (with a closed cap) at 37°C for 48 h. Identification of the cultured bacteria was carried out using the Vitek 2 compact system (BioMérieux). The biochemical patterns for two bacterial isolates were received with the use of the ANC ID card of the Vitek 2 compact system (BioMérieux).

In view of the fact that the same bacterial species, *C. perfringens*, was isolated from both patients and because of its pathogenic potential and epidemiological consequences, the following strains of *C. perfringens*: from patient No. 1 – isolate no. 8554/M/17; from patient No. 2 – isolate no. 8567/M/17, were preserved and transferred to the Chair of Microbiology, Jagiellonian University Medical College, Krakow, Poland for further studies. The *C. perfringens* strain cultured from the specimen obtained from a rectal swab from patient No. 2 was not preserved, which made it impossible to conduct further analyses for this strain.

The strains 8554/M/17 and 8567/M/17 were stored with the use of Cryobank (BioMaxima) at -70°C. *C. perfringens* 3624 ATCC (The American Type Culture Collection) standard was used as a reference strain.

Antibiotic susceptibility testing. To determine the drug-resistance profiles, the E-test method was used, enabling determination of MIC (Minimal Inhibitory Concentration) for: amoxicillin – AML, penicillin – P,

Table I
Antibiotic susceptibility for *C. perfringens* isolates (The E-test method, according to EUCAST 2017).

Isolate no.	Penicillins			Cephalosporins			Carbapenems			Fluoroquinolones	Glycopeptide and lipoglycopeptides		Macrolides, incosamides		Tetracyclines	Miscellaneous agents		
	AML	P	PRL	XL	CRO	CTX	DOR	ETP	IMP		MEM	TEC	VA	E		DA	TE	C
8554/M/17 (Patient 1)	MIC mg/l	0.016	0.016	0.016	0.016	0.016	0.016	0.016	0.016	0.016	0.047	0.50	0.50	0.50	4	0.19	0.016	0.032
	S/R	S	S	S	S	S	S	S	S	S	S	S	R	S	R	S	S	S
8567/M/17 (Patient 2)	MIC mg/l	0.016	0.016	0.016	0.016	0.016	0.016	0.016	0.016	0.016	0.047	0.50	0.50	0.50	4	0.125	0.016	0.032
	S/R	S	S	S	S	S	S	S	S	S	S	S	R	S	R	S	S	S

AML – amoxicillin, P – penicillin, PRL – piperacillin, XL – amoxicillin/clavulanic acid, CRO – ceftriaxone, CTX – cefotaxime, DOR – doripenem, ETP – ertapenem, IMP – imipenem, MEM – meropenem, MXF – moxifloxacin, TEC – teicoplanin, VA – vancomycin, E – erythromycin, DA – clindamycin, TE – tetracycline, C – chloramphenicol, M – metronidazole, RD – rifampicin, MIC – Minimal Inhibitory Concentration, S/R – susceptibility, R – resistant, S – susceptible

piperacillin – PRL, amoxicillin/clavulanic acid – XL, ceftriaxone – CRO, cefotaxime – CTX, doripenem – DOR, ertapenem – ETP, imipenem – IMP, meropenem – MEM, moxifloxacin – MXF, erythromycin – E, clindamycin – DA, tetracycline – TE, chloramphenicol – C, metronidazole – M, rifampicin – RD and with Glycopeptide Resistance Detection (GRD) for: teicoplanin – TEC and vancomycin – VA. The results were interpreted according to The European Committee on Antimicrobial Susceptibility Testing (EUCAST 2017) – Table I.

PCR multiplex. To isolate DNA, the Genomic Mini Set (A&A Biotechnology) was used according to the manufacturer's protocol. The presence of the genes encoding toxins of *C. perfringens* was confirmed using multiplex PCR amplification according to van Asten et al. (2009) with specific primers (Genomed). The following fragments of the genes were detected (the gene product, length of the fragment): *cpa* (α -toxin, 324 bp); *cpb* (β -toxin, 195 bp); *cpb2* (β 2-toxin, 548 bp); *etx* (ϵ -toxin, 376 bp); *iap* (ζ -toxin, 272 bp); *cpe* (enterotoxin, 485 bp). The final images from electrophoresis were processed using QuantityOne software, as well as GelDoc2000 device (Bio-Rad, USA).

Molecular typing with PFGE. The chosen *C. perfringens* isolates underwent molecular typing using the PFGE method according to the methodology described by Maslanka et al. (1999). Chromosomal DNA of the bacterial strains was isolated in agarose blocks and then digested with the use of restriction enzyme *Sma*I (MBI Fermentas). Electrophoretic separation was performed with the CHEF-DR II (Bio-Rad) instrument and restriction analysis was carried out using the Gel-Compar II (Applied Maths) software with the application of UPGMA clustering method and Jaccard index. The obtained genetic profiles were interpreted according to the guidelines given by van Belkum et al. (2007).

The profiles of the strains under investigation that were obtained using PCR multiplex and PFGE were compared with restriction patterns of the strains that came from the event in 2015, when in the Orthopedics and Traumatology Department of the same hospital, three cases of gas gangrene caused by *C. perfringens* were detected simultaneously (Brzychczy-Włoch et al. 2016).

Results

The two *C. perfringens* isolates studied (8554/M/17 from patient No. 1 and 8567/M/17 from patient No. 2) had identical biochemical profiles. Based on the results obtained using E-test, the same pattern of antibiotic susceptibility of the strains from the two examined patients was demonstrated (Table I).

Multiplex PCR confirmed the presence of the *cpa* gene encoding α -toxin for both *C. perfringens* isolates studied. Moreover, both strains demonstrated the presence of the *cpb2* gene encoding β 2-toxin (Fig. 1). A comparison of the results from 2017 and the genetic analysis of the strains from the event in 2015 enabled us to demonstrate that the currently examined strains have the genes encoding the same toxins (α and β 2) as the two strains analyzed in 2015: no. 7143 (patient No. 3) and no. 7149 (patient No. 2) (Brzychczy-Włoch et al. 2016).

As a result of the molecular analysis conducted using PFGE, two genetically different, unique restriction profiles were found corresponding to two different clones of the *C. perfringens* isolates studied in 2017: isolate no. 8554/M/17 – clone A; isolate no. 8567/M/17 – clone B (Fig. 2A).

Profiles of the strains under investigation that were obtained using PFGE were then compared with restriction patterns of the strains from the event in 2015 (Brzychczy-Włoch et al. 2016). A comparative analysis of PFGE profiles from 2017 and the results from the event from 2015 found one identical and genetically unique restriction profile, corresponding to one clone of *C. perfringens* for two strains: no. 8567/M/17 (patient No. 2 in 2017) and no. 7143 (patient No. 3 in 2015) (Fig. 2B).




Fig. 1. Detection of genes encoding virulence factors of *C. perfringens* isolates in 2017 tested by multiplex PCR.

Legend: M – size marker,

1 – *C. perfringens* isolate no. 8567/M/17 from patient no. 2

2 – *C. perfringens* isolate no. 8554/M/17 from patient no. 1

3 – reference strain of *C. perfringens* ATCC 3624

cpb2 – gene of β 2-toxin; *cpa* – gene of α -toxin




Fig. 2. Genetic profiles of *C. perfringens* isolates, subjected to DNA digestion by restriction enzyme *Sma*I with the use of pulsed-field gel electrophoresis (PFGE) and GelCompar II software.

Legend: A – Comparison of two isolates from 2017; B – comparison of five isolates: three from 2015 and two from 2017
 8554/M/17 – *C. perfringens* isolate isolated from patient no. 1 in 2017; ATCC 3624 – reference strain of *C. perfringens*
 8567/M/17 – *C. perfringens* isolate isolated from patient no. 2 in 2017
 7143 – *C. perfringens* isolate isolated from patient no. 3 in 2015; 7149 – *C. perfringens* isolate isolated from patient no. 2 in 2015
 7013 – *C. perfringens* isolate isolated from patient no. 1 in 2015

Due to this fact, an attempt was undertaken to find a connection between patient No. 2 from 2017 and patient No. 3 from 2015. In the course of an epidemiological investigation, it was only found that both patients were hospitalized in the same hospital, but they were in different wards at different times. The patient from whom the strain no. 7143 was isolated (patient No. 3) was hospitalized in April 2015 at the Department of Orthopedics and Traumatology. The currently described patient No. 2 from whom strain no. 8567/M/17 was isolated in 2017 was in the General Surgery Department in May 2014. Both patients underwent surgery in the main operating block of the hospital. Kinship, close contact, possession of the same animals and residence in each other's neighborhood were excluded.

Discussion

The cases presented above illustrate the clinical picture of necrotizing soft tissue infection, caused by *C. perfringens* toxinotype A. The infection leading to amputation of the left lower limb in the patient No. 1

makes us to assume that he developed type 3 NSTI. Atherosclerosis of the lower extremities, alcoholism, and neuropathy with subsequent vascular insufficiency contributed to the rapid progression of the disease. The course of the infection in patient No. 2, which was classified as Fournier gangrene and isolation of the etiologic agent from the wound, i.e. *C. perfringens*, as well as medical history pointing to decreased immunity and advanced neoplastic disease of the large intestine, allowed to confirm this infection as type 1 NSTI.

The results of microbiological testing with phenotypic methods (biochemical patterns and antibiotic susceptibility testing) did not demonstrate differences between the strains isolated from biological specimens from the patients. On the other hand, the application of molecular methods allowed to characterize each of them accurately and compare them with strains isolated during the epidemiological investigation, which took place in the same hospital during the event in 2015 (Brzychczy-Włoch et al. 2016).

Determination of the toxinotype of *C. perfringens* isolates was possible owing to the use of multiplex PCR. The strains identified for both patients described

in this study were type A and had the genes encoding α -toxin (*cpa* gene) and β 2-toxin (*cpb2* gene). The role of α -toxin, which is the main virulence factor in of *C. perfringens* in gas gangrene, seems to be well known and is chiefly associated with hemolysis and dermonecrosis. To the best of our knowledge, there is no evidence pointing to β 2-toxin's contribution to necrotizing soft tissue infections in humans. Despite this, in the work reporting three simultaneous cases of gas gangrene associated with *C. perfringens* type A strains in 2015, two patients carried the strains that had the *cpb2* gene, indicated an extremely severe course of these infections (Brzychczy-Włoch et al. 2016). The presence of the gene for β 2-toxin in *C. perfringens* strains causing soft tissue infections in humans requires further observation, which could be assisted by the application of genetic analysis of this pathogen in every clinical case involving *C. perfringens*.

Molecular diagnostics of the strains isolated also allowed their final differentiation and an attempt to determine their origin. Owing to macrorestriction analysis of chromosomal DNA combined with PFGE, it was possible to determine that the strain of the described patient No. 1 turned out to be different from the strains isolated from the patient No. 2. However, the identification of an identical restriction profile for the *C. perfringens* isolate from patient No. 2 and the profile of the strain isolated from one of the patients during the event in 2015 in the same hospital (isolate no. 7143 from patient No. 3) deserves more attention. According to the literature, the species *C. perfringens* is characterized by remarkable diversity and numerous mutations due to the presence of genes encoding toxins not only in the chromosome but also in plasmids (Myers et al. 2006; Park et al. 2016; Kiu et al. 2017). Hence, a random detection of the presence of an identical strain in two distant, independent events is unlikely. However, a high genetic similarity between *C. perfringens* strains can be demonstrated when they cause epidemiologically related infections (Johansson et al. 2006; Xiao et al. 2012). Our epidemiological analysis points to the fact that patient No. 2 (from the event in 2017) had already been subjected to a surgical procedure within the abdominal cavity in this hospital in 2014. There is a possibility that he had already been a carrier of *C. perfringens* in the gastrointestinal tract at that time and that he became the source of the infection and as result of the surgical procedure the bacterium appeared in the operating block of the hospital. It is known that *C. perfringens* may be present everywhere, even in dust, and is capable of producing spores, which can survive in unfavorable conditions for many months (van Asten et al. 2010; Kędzińska et al. 2012; Brzychczy-Włoch and Bulanda 2014). In 2014, there was no reason for routine use of sporicidal substances in the operating block (no

symptomatic cases of infection with the bacillus *C. perfringens* at the hospital). One should, then, take into consideration the possibility of survival of *C. perfringens* spores in the hospital after the hospitalization of patient No. 2 in 2014 and the possible infection of one of the three patients undergoing surgery in the same operating block in 2015. Even more, other connections between both patients were excluded (kinship or being neighbors). During the event in 2015, the appropriate epidemiological and protective procedures against an outbreak were implemented (including the application of sporicidal agents). Therefore, it is unlikely that the *C. perfringens* strain survived two more years in the operating block (Brzychczy-Włoch et al. 2016). Additionally, the result of rectal swab testing in patient No. 2 in 2017 indicated that the isolate from that sample is an endogenous (own) strain of the patient. It was patient No. 2 who probably had been a carrier of the described *C. perfringens* strain in the gastrointestinal tract for years and his decreased immunity together with cancer created the favorable conditions for the growth of bacteria and development of infection in 2017 (Brzychczy-Włoch and Bulanda 2014; Kuzaka et al. 2018).

Our research has some limitation as there are no environmental or hospital staff studies that could have precisely demonstrated the source where *C. perfringens* clone that was able to survive. However, owing to the application of molecular methods, it was possible to determine that the patients simultaneously hospitalized in 2017, in a single Department of General Surgery developed two different types of NSTI caused by two different *C. perfringens* clones. Archival data from 2015 allowed the identification of an identical clone from the same hospital. The employment of genetic analyses also enabled us to document, in the strains studied in 2015 and in 2017, the presence of the *cpb2* gene encoding β 2-toxin. Moreover, the epidemiological data and detailed analysis of the course of both events in the hospital made it possible to attempt to understand the reasons of the survival of the *C. perfringens* clone in the operating block and suggest that this bacterium may have survived in adverse conditions of the external environment for many months, posing a potential threat to patients. Hence, compliance with the procedures concerning the operating block hygiene must always be strictly observed. Maybe it is also worthwhile to consider modifying them and to introduce periodic (e.g. every three months) mandatory application of sporicidal agents regardless of whether there were clinical cases of *C. perfringens* infection or not.

ORCID

Dominika Salamon [0000-0002-5974-5520](https://orcid.org/0000-0002-5974-5520)

Małgorzata Bulanda [0000-0002-4679-9263](https://orcid.org/0000-0002-4679-9263)

Monika Brzychczy-Włoch [0000-0002-7415-0154](https://orcid.org/0000-0002-7415-0154)

Acknowledgments

The authors would like to thank the employees of the General Surgery Department of the Provincial Specialist Hospital in Włocławek for their cooperation in conducting the study. Language translation: Catherine Ridgetile.

Conflict of interest

The authors do not report any financial or personal connections with other persons or organizations, which might negatively affect the contents of this publication and/or claim authorship rights to this publication.

Literature

- Allaart JG, van Asten AJAM, Vernooij JCM, Gröne A. Beta2 toxin is not involved in *in vitro* cell cytotoxicity caused by human and porcine cpb2-harboring *Clostridium perfringens*. *Vet Microbiol*. 2014;171(1–2):132–138. <https://doi.org/10.1016/j.vetmic.2014.03.020>
- Brzywczy-Włoch M, Bulanda M. Analysis of genetic similarities between *Clostridium perfringens* isolates isolated from patients with gas gangrene and from hospital environment conducted with the use of the PFGE method. *Pol J Surg*. 2014;86(3):141–146. <https://doi.org/10.2478/pjs-2014-0026>
- Brzywczy-Włoch M, Ochońska D, Piotrowska A, Bulanda M. Gas gangrene of different origin associated with *Clostridium perfringens* type A in three patients simultaneously hospitalized in a single department of Orthopedics and Traumatology in Poland. *Pol J Microbiol*. 2016;65(4):399–406. <https://doi.org/10.5604/17331331.1227665>
- Johansson A, Aspan A, Bagge E, Båverud V, Engström BE, Johansson KE. Genetic diversity of *Clostridium perfringens* type A isolates from animals, food poisoning outbreaks and sludge. *BMC Microbiol*. 2006;6(1):47. <https://doi.org/10.1186/1471-2180-6-47>
- Kędzińska J, Obuch-Woszczyński P, Pituch H, Młynarczyk G. [*Clostridium perfringens* as the etiological agent of antibiotic associated diarrhea] (in Polish). *Postepy Mikrobiol*. 2012;51(1):17–25.
- Kiu R, Caim S, Alexander S, Pachori P, Hall LJ. Probing genomic aspects of the multi-host pathogen *Clostridium perfringens* reveals significant pangenome diversity, and a diverse array of virulence factors. *Front Microbiol*. 2017;8:2485. <https://doi.org/10.3389/fmicb.2017.02485>
- Kuzaka B, Wróblewska MM, Borkowski T, Kawecki D, Kuzaka P, Młynarczyk G, Radziszewski P. Fournier's gangrene: clinical presentation of 13 cases. *Med Sci Monit*. 2018;24:548–555. <https://doi.org/10.12659/MSM.905836>
- Maslanka SE, Kerr JG, Williams G, Barbaree JM, Carson LA, Miller JM, Swaminathan B. Molecular subtyping of *Clostridium perfringens* by pulsed-field gel electrophoresis to facilitate food-borne-disease outbreak investigations. *J Clin Microbiol*. 1999;37(7):2209–2214.
- Myers GSA, Rasko DA, Cheung JK, Ravel J, Seshadri R, DeBoy RT, Ren Q, Varga J, Awad MM, Brinkac LM, et al. Skewed genomic variability in strains of the toxigenic bacterial pathogen, *Clostridium perfringens*. *Genome Res*. 2006;16(8):1031–1040. <https://doi.org/10.1101/gr.5238106>
- Navarro M, McClane B, Uzal F. Mechanisms of action and cell death associated with *Clostridium perfringens* toxins. *Toxins (Basel)*. 2018;10(5):212. <https://doi.org/10.3390/toxins10050212>
- Park M, Deck J, Foley SL, Nayak R, Songer JG, Seibel JR, Khan SA, Rooney AP, Hecht DW, Rafii F. Diversity of *Clostridium perfringens* isolates from various sources and prevalence of conjugative plasmids. *Anaerobe*. 2016;38:25–35. <https://doi.org/10.1016/j.anaerobe.2015.11.003>
- Rood JI, Adams V, Lacey J, Lyras D, McClane BA, Melville SB, Moore RJ, Popoff MR, Sarker MR, Songer JG, et al. Expansion of the *Clostridium perfringens* toxin-based typing scheme. *Anaerobe*. 2018;53:5–10. <https://doi.org/10.1016/j.anaerobe.2018.04.011>
- Stevens DL, Aldape MJ, Bryant AE. Life-threatening clostridial infections. *Anaerobe*. 2012;18(2):254–259. <https://doi.org/10.1016/j.anaerobe.2011.11.001>
- Stevens DL, Bryant AE. Necrotizing Soft-Tissue Infections. *N Engl J Med*. 2017;377(23):2253–2265. <https://doi.org/10.1056/NEJMra1600673>
- EUCAST. Breakpoint tables for interpretation of MICs and zone diameters Ver. 7.1 [Internet]. Basel (Switzerland): European Committee on Antimicrobial Susceptibility Testing; 2017 [cited 2019 Jan 21]. Available from http://www.eucast.org/ast_of_bacteria/previous_versions_of_documents/
- van Belkum A, Tassios PT, Dijkshoorn L, Haegman S, Cookson B, Fry NK, Fussing V, Green J, Feil E, Gerner-Smidt P, et al.; European Society of Clinical Microbiology and Infectious Diseases (ESCMID) Study Group on Epidemiological Markers (ESGEM). Guidelines for the validation and application of typing methods for use in bacterial epidemiology. *Clin Microbiol Infect*. 2007;13 Suppl 3:1–46. <https://doi.org/10.1111/j.1469-0691.2007.01786.x>
- van Asten AJAM, Nikolaou GN, Gröne A. The occurrence of cpb2-toxigenic *Clostridium perfringens* and the possible role of the β 2-toxin in enteric disease of domestic animals, wild animals and humans. *Vet J*. 2010;183(2):135–140. <https://doi.org/10.1016/j.tvjl.2008.11.005>
- Wróblewska M, Kuzaka B, Borkowski T, Kuzaka P, Kawecki D, Radziszewski P. Fournier's gangrene – current concepts. *Pol J Microbiol*. 2014;63(3):267–273.
- Xiao Y, Wagendorp A, Moezelaar R, Abee T, Wells-Bennik MHJ. A wide variety of *Clostridium perfringens* type A food-borne isolates that carry a chromosomal *cpe* gene belong to one multilocus sequence typing cluster. *Appl Environ Microbiol*. 2012;78(19):7060–7068. <https://doi.org/10.1128/AEM.01486-12>
- Żaba R, Grzybowski A, Prokop J, Żaba Z, Żaba C. Fournier's gangrene: historical survey, current status, and case description. *Med Sci Monit*. 2009;15(2):CS34–CS39.