

# Terrestrial Non-Parasitic Nematode Assemblages associated With Glyphosate-tolerant and Conventional Soybean-Based Cropping Systems

Akhona Mbatyoti,<sup>1,2\*</sup> Mieke Stefanie Daneel,<sup>2</sup> Antoinette Swart,<sup>3</sup> Dirk de Waele,<sup>1,4</sup> and Hendrika Fourie<sup>1</sup>

<sup>1</sup>Unit for Environmental Sciences and Management, Potchefstroom Campus, North-West University, Potchefstroom, South Africa.

<sup>2</sup>Agricultural Research Council–Tropical and Subtropical Crops, Nelspruit, South Africa.

<sup>3</sup>Agricultural Research Council–Plant Health and Protection, Pretoria, South Africa.

<sup>4</sup>Laboratory of Tropical Crop Improvement, Faculty of Bioscience Engineering, Department of Biosystems, University of Leuven, Willem de Croylaan 42, 3001 Leuven, Belgium.

\*E-mail: MbatyotiO@arc.agric.za.

This paper was edited by Eyuaem Abebe.

Received for publication January 7, 2018.

## Abstract

Information about the effects of glyphosate on nematodes is limited and contradictory, while none existing for South African agricultural fields. The abundance and identity of non-parasitic nematodes in the rhizospheres of commercial glyphosate-tolerant and conventional (non-glyphosate-tolerant), soybean cultivars from cultivated fields, and adjacent natural vegetation (reference system) were obtained for two growing seasons. The impact of glyphosate was also investigated on non-parasitic nematodes in a 2-year soybean-maize cropping system. Thirty-two non-parasitic nematode genera were identified from soils of the three field ecosystems, with most of the genera occurring in natural vegetation (28), and less in conventional (23) and glyphosate-tolerant soybean (21). Bacterivores had the greatest diversity in soils of all three ecosystems during both seasons, while fungivores tended to be more abundant in glyphosate-tolerant soybean fields especially during the second season. Soils from the three ecosystems were disturbed and degraded with low abundance and diversity of omnivores and predators. Of the 14 genera identified from the soybean-maize cropping experiment, bacterivores dominated in terms of diversity in non-treated, and fungivores in glyphosate-treated plots. Soils from glyphosate-treated plots were degraded, less enriched and fungal-mediated, while those from non-treated plots were disturbed, enriched, and bacterial-mediated.

## Key words

Assemblages, Non-parasitic nematodes, Soybean.

Commercial production of genetically modified crops (GM), either herbicide or insect tolerant, commenced in the 1990s (Dill et al., 2008; Shütte et al., 2017). Herbicide tolerance of GM crops to broad spectrum herbicides containing glyphosate (N-phosphonomethyl) as the active substance is the predominant trait of these crops (Newman et al., 2016). Among glyphosate tolerant crops grown globally, soybean (*Glycine max* L.) dominates in terms of hectares planted (54.2 million ha), followed by maize (*Zea mays* L.) (13.2 million ha), cotton (*Gossypium hirsutum* L.) (5.1 million ha) and canola (*Brassica napus* L.) (2.3 million ha) (Shütte et al., 2017). In South Africa, glyphosate became commercially available more than a decade ago and is now widely

used in soybean- and maize-based cropping systems in particular (Dlamini et al., 2014). It is estimated that more than 90% of soybean (630,000 ha) and 16% of maize (284,000 ha) grown in South Africa are glyphosate tolerant (Dlamini et al., 2014; James 2015). The driving force behind the rapid adoption of glyphosate is because producers prefer to use a single herbicide to control a broad spectrum of weeds and grasses, resulting in minimal crop injury and great economic benefits to producers (Hurley et al., 2009).

Glyphosate is often regarded as an environmentally-friendly pesticide due to its low mammalian toxicity, relatively short environmental half-life and very low activity in soil due to its binding to soil minerals (Duke

and Powles, 2008; Cerdeira and Duke, 2010). However, the increasing cultivation of glyphosate tolerant crops has raised a wide range of concerns such as its effects on non-target micro-organisms, e.g., nematodes in the soil (Zhao et al., 2013; Allegrini et al., 2015; Newman et al., 2016). Nematodes play a crucial role in important ecosystem services such as nutrient recycling and decomposition, suppression of pathogenic micro-organisms, and biodegradation of harmful compounds (Bongers and Bongers, 1998; Ferris et al., 1998; Neher, 2001; Wardle et al., 2005). As a result, changes in nematode community composition (assemblage) may have a substantial impact on the ecosystem functioning (Wada et al., 2011; Fraschetti et al., 2016).

Information about the non-target effects of glyphosate on soil nematodes is scarce and not well documented. More important, often inconclusive and/or conflicting effects of glyphosate on nematode assemblages are reported. Only six scientific reports could be found that dealt with the effects of glyphosate on nematodes. The majority of these focused on the effects that glyphosate has on plant-parasitic nematodes (Osman and Viglierchio, 1981; Vega et al., 1993; Yang et al., 2002, Liphadzi et al., 2005; Cerdeira et al., 2007; Noel and Wax, 2009). Liphadzi et al. (2005), however, reported that different glyphosate

dosages had no effect on non-parasitic nematode densities in a growth chamber experiment.

No information on the effects of glyphosate on, or its association with either plant-parasitic or terrestrial non-parasitic nematodes (generally referred to as beneficial or free-living), is available for South African agricultural production areas. Therefore, the main aims of this study were to (i) identify terrestrial, non-parasitic nematode assemblages in commercial soybean fields where glyphosate has been applied regularly versus not applied for at least 5 years prior to this study and (ii) examine whether glyphosate application affected such nematode assemblages in a 2 year soybean-maize cropping system.

## Materials and methods

### Commercial soybean field study

During the 2011/12 growing season, rhizosphere soil was collected from soybean plants that were cultivated at eight local fields. Four of these fields were planted with glyphosate-tolerant and four with conventional soybean cultivars (Fig. 1), representing the two soybean ecosystems. Concurrently, soil samples were also collected from a third ecosystem, viz. natural vegetation




Figure 1: Location of the six localities where terrestrial, non-parasitic nematodes were sampled from soybean fields (2011/12: red triangles and 2012/13: blue triangles) and adjacent natural vegetation (yellow triangles) during two consecutive growing seasons (Map compiled by: Ms L. de Swart, NWU).

(representing a reference system) either adjacent to, or within 50 to 100m from the soybean fields sampled. From each of these three ecosystems, at each sampling locality 80 rhizosphere soil samples were collected, pooled and 20 sub-samples examined.

Glyphosate had been applied continuously for a minimum of 5 years prior to our study in the fields where glyphosate-tolerant soybean and/or maize cultivars were cultivated. However, in the fields planted with conventional soybean no glyphosate-tolerant cultivars were grown and no glyphosate applied for at least 5 years prior this study or never before. No crop cultivation has taken place for at least 10 years prior to this study in the areas where the natural vegetation was sampled.

During the 2012/13 growing season, the same fields sampled during the preceding season were sampled again as well as nine additional fields and adjacent nat-

ural vegetation (Fig. 1). Five of these additional fields were planted with glyphosate-tolerant and four with conventional soybean cultivars, with information about the soybean cultivar planted, crop history and soil properties for each field sampled being supplied in Table 1. Soil properties for each site were determined by the EcoAnalitica Laboratory of North-West University (NWU, Potchefstroom) using internationally-accredited protocols (Walkey and Black, 1947; Bouyoucos, 1962; Beretta et al., 2014). Mean rainfall and temperature data, obtained from the database of the Agricultural Research Council, Institute for Soil, Climate and Water, AgroClimatology for each site, from planting of the soybean crops until nematode sampling are also listed (Table 2). Rip and till was the soil cultivation practice used in all soybean fields sampled.

Nematodes were extracted from 200g soil samples using the decanting and sieving method (Hooper

**Table 1. Soybean cultivar planted at each soybean field, crop history, and selected soil chemical and physical properties of each field where plant-parasitic nematodes from roots and rhizosphere soil samples were collected during the 2011/12 and 2012/13 growing seasons.**

Sampling season	Locality	Ecosystem and cultivar	Crop history	pH (H <sub>2</sub> O)	Soil chemical properties					Soil physical properties			
					Ca	Mg	K	Na	P	% Sand	% Silt	% Clay	% total C
2011/12 and 2012/13	Bothaville	Glyphosate-tolerant soybean (PAN1664R)	Maize/Sunflower	6.48	381	107	205	0.5	204	94.7	0.7	4.6	0.21
2012/13		Glyphosate-tolerant soybean (PAN1664R)	Maize/Sunflower	6.48	437	81	170	0.5	170	94.5	0.7	4.8	0.10
2011/12 and 2012/13		Conventional soybean (Egret)	Maize/Sunflower	6.89	581	81	246	1	166	92.6	0.7	6.7	0.23
2012/13		Conventional soybean (Egret)	Maize/Sunflower	6.43	374	96	203	0.5	203	94.6	0.7	4.7	0.22
2011/12 and 2012/13		Natural vegetation (grass)		6.77	446	78	194	0.5	169	94.5	0.7	4.7	0.21
2011/12 and 2012/13		Natural vegetation (grass)		6.06	574	165	400	5	252	89.6	3.4	7	1.5

## Non-parasitic nematodes of grains

2011/12 and 2012/13	Brits	Glyphosate- tolerant soybean (PAN1583R)	Maize/ Sunflower	7.28	1,434	304	497	2	500	79.9	5.5	14.6	1.36
2011/12 and 2012/13		Conventional soybean (Egret)	Soybean/ Wheat	7.49	1,699	346	291	58.5	399.	81.4	7.6	11	0.76
2011/12 and 2012/13		Natural vegetation (grass)		7.11	2,840	559	417	91.5	509	58.3	12.4	29.2	1.51
2011/12 and 2012/13		Natural vegetation (grass)		7.6	3,206	636	342	98	53	74.1	9.1	16.8	3.95
2012/13	Edenville	Glyphosate- tolerant soybean (PAN1664R)	Soybean	6.15	1,486	401	419	60.5	40	59.3	19.1	21.6	0.20
2012/13		Conventional soybean (Superboon)	Soybean	4.97	512	213	292	9.5	118	70.7	9.3	20	0.49
2012/13		Natural vegetation (grass)		5.92	396	67	268	3	93	85.9	3.4	10.7	3.89
2012/13		Natural vegetation (grass)		5.64	222	52	271	1.5	86	86.6	3.2	10.1	0.34
2011/12 and 2012/13	Marble Hall	Glyphosate- tolerant soybean (PAN1454R)	Maize/ Soybean	6.09	666	228	390	20	402	84.9	6.7	8.4	0.56
2012/13		Glyphosate- tolerant soybean (LS6164R)	Maize/ Soybean	6.64	541	149	146	23	118	91	3.7	5.1	0.84
2012/13		Glyphosate- tolerant soybean (LS6164R)	Maize/ Soybean	6.07	826	212	291	176	336	88.7	3.7	7.6	0.40
2011/12 and 2012/13		Conventional soybean (Egret)	Soybean	7.05	1,001	402	390	34.5	67	78.6	8.7	12.7	0.58
2012/13		Conventional soybean (MC555)	Soybean	6.62	1,012	244	509	14.5	156	91.2	3.7	5.1	1.76
2011/12 and 2012/13		Natural vegetation (grass)		6.83	968	238	346	35.5	151	69.2	9.3	21.5	1.8
2011/12 and 2012/13		Natural vegetation (grass)		6.52	455	104	192	24.5	164	86.1	1.4	12.6	0.54
2011/12 and 2012/13		Natural vegetation (grass)		5.93	810	121	375	25	419	83.7	3.8	12.5	2.54

2012/13	Viljoenskroon	Glyphosate-tolerant soybean (PAN1583R)	Maize/Soybean	6.82	381	96	330	5	295	94.1	1.2	4.7	0.2
2012/13		Natural vegetation (grass)		6.14	433	195	370	9.5	227	84.4	5.9	9.7	1.14
2011/12 and 2012/13	Winterton	Glyphosate-tolerant soybean (PAN6164R)	Soybean	5.67	1,782	400.5	381	20.5	85	81.2	10.4	8.5	2.49
2012/13		Conventional soybean (Mukwa)	Maize/Soybean	6.33	1,565	198	182	2	390	78.8	12.9	8.4	1.3
2011/12 and 2012/13		Conventional soybean (Mukwa)	Maize/Soybean	5.92	1,827	139	551	5.5	532	44.8	38.4	16.7	2.77
2011/12 and 2012/13		Natural vegetation (grass)		6.85	751	120	86	0.5	335	61.4	23.3	15.3	1.62
2011/12 and 2012/13		Natural vegetation (grass)		5.65	887	208	295	4.5	358	45.9	31.4	22.8	2.59

et al., 2005), and counted and identified to genus level using a 1-ml Hawksley slide and light microscope (1,000 × magnification) (Doncaster et al., 1967). This process was repeated once for each sample and the

mean of the two counts were used for data analyses. At least 30 individuals from each genus per sample were, after counting, fixed in a heated formaldehyde-propionic-acid-water (FPG) solution (100ml of

**Table 2. Average temperature and rainfall figures for the 28 sites where a nematode survey was conducted during the 2011/12 and 2012/13 growing seasons.**

Locality and province	Growing season	Temperature (°C)		Rainfall (mm)
		Min.	Max.	
Bothaville (Free State)	2011/12	13	26	272
	2012/13	14	30	255
Brits (North West)	2011/12	16	33	414
	2012/13	16	32	365
Edenville (Free State)	2012/13	16	32	303
Marble Hall (Mpumalanga)	2011/12	18	31	402
	2012/13	18	35	353
Viljoenskroon (Free State)	2012/13	14	31	373
Winterton (Kwa-Zulu Natal)	2011/12	14	30	409
	2012/13	14	30	417

a 40% formalin solution, 10ml propionic acid and 890ml distilled water). The glass dish with the fixed nematodes were placed in an incubator at 40°C for 72hr and the FPG solution stepwise replaced with glycerin (Marais et al., 2017). The fixed nematodes were hand-picked from the glycerin using a fine-tip needle and permanently mounted in glycerin on glass microscope slides according to the paraffin-ring protocol (Hooper, 1986). Genus identification of nematodes was done and verified by Dr Antoinette Swart, a nematode specialist-taxonomist of the Agricultural Research Council – Plant Health and Protection (Roodeplaat, South Africa).

### Soybean-maize cropping experiment

The experimental site consisted of a small field (0.028ha plot) situated on the premises of the Agricultural Research Council's Grain Crops Institute, Potchefstroom, South Africa. The study was conducted over two consecutive growing seasons (2013/14 and 2014/15) with soybean being cultivated during the first and maize during the second season. The soil of the plot contained 94% sand and 6% clay. The organic matter content ranged from 0.18% (2013/14 season) to 0.23% (2014/15 season), while soil pH (H<sub>2</sub>O) was 8 for the 2013/14 season and 7.8 for the 2014/15 season. The history about crops grown and herbicides applied on the experimental site, glyphosate application dosages and dates during the experimental period, nematode sampling dates and rainfall, and temperature data are supplied (Table 3).

The experimental plot was split into two halves (0.013ha each), which were divided by a fallow, 2-m buffer strip. Before planting the plot for the first season with soybean, weeds that grew on the experimental plot were mechanically hoed and left on the experimental plot. This is the practice that local farmers use. On 18 November 2013, at the beginning of the 2013/14 growing season, seeds of the glyphosate-resistant soybean cultivar LS 6164R were planted after the soil was ripped and tilled using a tractor. The soybean seeds were planted (170 per row) in 5-m-long rows with intra- and inter-row spacings of 3cm and 0.9m, respectively. Each seed was coated with *Bradyrhizobium japonicum* race WB74 at the recommended dosage rate (Soygro Pty Ltd; www.soygro.co.za). The layout of the experiment was a split-plot design with 12 replicates. Each row represented a replicate.

After germination, soybean seedlings were irrigated with ~25mm water three times a week using a sprinkler irrigation system, except when it had rained sufficiently. When naturally occurring weeds were 10 to 20cm tall, one half of the experimental plot was

treated with glyphosate (active substance 360g/l glyphosate present as 441g/l of the potassium salt at a dosage rate of 2l/ha) using a knapsack sprayer. Applications were done early in the mornings to avoid wind and possible drift of the product as specified by the owner company of the product used. The other half of the experimental plot was not treated with glyphosate or any other herbicide and represented the control. Weeds in the non-treated plot were removed using a hand hoe and left on the soil. This meant that the upper surface of the soil was disturbed during the hoeing action and organic material was left on the soil to decompose.

Before planting, as well as 10 to 20 days after each glyphosate application and also at 120 to 140 days after planting (i.e., at crop maturity), rhizosphere soil and the root systems of nine soybean plants from each replicate were collected, thoroughly mixed and one sub-sample per replicate used for nematode analyses.

During the winter of 2014, no crop was grown and both halves of the experimental plot were left fallow without any weed control being applied. However, before planting seeds of the glyphosate-resistant maize cultivar DKC 80–30 RR on 18 November 2014 of the follow-up growing season, glyphosate was applied on the same plot half where glyphosate had been applied during the previous season (where soybean was planted). Again, the other half of the experimental plot was not treated and the weeds hand hoed and left on the soil. Ten days later, the soil was ripped and tilled and seeds of commercially available maize cultivar DKC 80–30 RR planted. Twenty-five maize seeds were planted per row, each being 5-m long, with intra- and inter-row spacings of 20cm and 0.9m, respectively. Two glyphosate applications were done as described above for the preceding soybean crop.

Ten to 20 days after each glyphosate application and also at 120 to 140 days after planting (i.e., at crop maturity), rhizosphere soil and the root systems of three maize plants from each replicate were collected, thoroughly mixed and one sub-sample per replicate used for nematode analyses. The same protocols were used for soil and root sampling, nematode extraction, counting, and identification to genus level as described for the commercial field sampling study.

### Data analyses

#### *Commercial soybean field study*

Nematode data were captured and  $\log_{10}(x+1)$  transformed using Microsoft Excel, Version 2013. Prominence values (PV) were calculated for each nematode genus using the protocol of De Waele and Jordaan

**Table 3. Crop history, agricultural practices, fertilisers applied, glyphosate application dates, and dosage rate, nematode sampling dates, rainfall, and minimum and maximum temperatures recorded of the experimental plot during the 2013/14 and 2014/15 growing seasons.**

Crop history: growing season, crop and herbicides applied	Growing season, crop and cultivar cultivated	Agricultural practice implemented	Inorganic fertiliser applied and dosage	Glyphosate application dates and dosage rates	Nematode sampling dates	Rainfall (mm)	Minimum temperature (°C)	Maximum temperature (°C)
2011/12, sunflower, no herbicide used (hand-hoeing of weeds)	2013/14 (1st year), soybean (cv. LS 6164 R)	Reap and plough before planting	None	10 January 2014 and 03 February 2014 @ 2 L/ha	1st = 30/01/2014; 2nd = 14/02/2014; 3rd = 22/04/2014	603	7.9	30.5
2012/13, maize, Gramoxone® (active substance bipyridyl 200 g/L as dichloride salt 276 g/L) dosage rate 3 L/ha	2014/15 (2nd year), maize (cv. DKC 80-30 RR)	Reap and plough before planting	2:3:2 (26) at planting @ 300 kg/ha Ureum @ 50 kg/ha 4 weeks after planting	17 December 2014 and 13 January 2015 @ 2 L/ha	1st = 08/01/2015; 2nd = 27/01/2015; 3rd = 11/03/2015	510	10.0	30.2

**Table 4. Non-parasitic nematodes associated with soybean and natural vegetation at 28 sites in the soybean production areas of South Africa during the 2011/12 and 2012/13 growing seasons (✓ indicates the presence of a genus; – indicates the absence of a genus).**

Genus	Functional guild <sup>a</sup> , followed by c–p value <sup>b</sup>	Glyphosate- tolerant soybean	Conventional soybean	Natural vegetation
<i>Mesorhabditis</i> (Osche, 1952); Dougherty, 1953	Ba1	✓	✓	✓
<i>Panagrolaimus</i> Fuchs, 1930	Ba1	✓	✓	✓
<i>Rhabditis</i> Dujardin, 1845	Ba1	✓	✓	✓
<i>Acrobeles</i> Linstow, 1877	Ba2	✓	✓	✓
<i>Acrobelloides</i> (Cobb, 1924); Thorne, 1937	Ba2	✓	✓	✓
<i>Cephalobus</i> Steiner, 1929	Ba2	✓	✓	✓
<i>Chiloplacus</i> Thorne, 1937	Ba2	✓	✓	✓
<i>Eucephalobus</i> Steiner, 1936	Ba2	✓	✓	✓
<i>Monhystera</i> Bastian, 1865	Ba2	✓	✓	✓
<i>Plectus</i> Bastian, 1865	Ba2	–	✓	✓
<i>Seleborca</i> Andrassy, 1985	Ba2	–	–	✓
<i>Wilsonema</i> Cobb, 1913	Ba2	–	–	✓
<i>Zeldia</i> Thorne, 1937	Ba2	–	–	✓
<i>Teratocephalus</i> de Man, 1876	Ba4	–	–	✓
<i>Alaimus</i> de Man, 1880	Ba4	–	✓	–
<i>Aphelenchoides</i> Fischer, 1894	Fu2	✓	✓	✓
<i>Aphelenchus</i> (Bastian, 1865); Cobb, 1927	Fu2	✓	✓	✓
<i>Ditylenchus</i> Filipjev, 1936	Fu2	✓	✓	✓
<i>Psilenchus</i> de Man, 1921	Fu2	✓	✓	✓
<i>Tylenchus</i> Bastian, 1865	Fu2	✓	✓	✓
<i>Coslenchus</i> Siddiqi, 1978	Fu3	–	–	✓
<i>Leptonchus</i> Cobb, 1920	Fu4	–	✓	✓
<i>Tylencholaimellus</i> (Cobb, 1915); de Man, 1921	Fu4	–	✓	✓
<i>Tylencholaimus</i> de Man, 1880	Fu4	✓	–	–
<i>Dorylaimus</i> Thorne, 1939	Om4	✓	✓	✓
<i>Eudorylaimus</i> Andrassy, 1959	Om4	✓	✓	✓
<i>Thornenema</i> Andrassy, 1959	Om4	✓	✓	✓



<i>Mononchus</i> Chitwood and Allen, 1959	Pr4	–	√	–
<i>Paraxonchium</i> Krall, 1958	Pr4	√	√	–
<i>Aporcelaimellus</i> Heyns, 1965	Pr5	√	–	√
<i>Discolaimium</i> Thorne, 1939	Pr5	√	–	√
<i>Discolaimoides</i> Heyns, 1963	Pr5	√	√	√

<sup>a</sup>Functional guilds (Ferris et al., 2001); <sup>b</sup>Colonizer-persister (c–p) values (Bongers, 1990). Trophic group with Ba, Bacterivores; Fu, Fungivores; Om, Omnivores; and Pr, Predators.

(1988).  $\log_{10}(x+1)$  transformed nematode data were also subjected to Student's *t*-test analyses using Statistica Version 13.2 (www.statsoft.com). This was done to determine whether any significant ( $P \leq 0.05$ ) differences existed between the predominant genera at each of the sampling sites with regard to the three ecosystems (viz. glyphosate-tolerant vs. conventional soybean, conventional soybean vs. natural vegetation and glyphosate-resistant vs. natural vegetation). The Mixed models analysis was also done using SPSS software (Version 25) to determine whether the three independent variables, e.g., season (2011/12 and 2012/13), location (eight for 2011/12 and 17 for 2012/13) and ecosystem (glyphosate-tolerant and conventional soybean, and natural vegetation), alone or interactively, affected the abundance of the various nematode trophic groups. In addition, nematode population density data were also illustrated on canonical correspondence analyses (CCA) triplots, using the Canoco 5 software package (www.canococ5.com). This way it was determined whether correlations existed for nematode genera and specific ecosystems for data, pooled across localities and per locality. Finally, to assess soil quality as expressed by the enrichment and structure values according to colonizer-persister (c–p) values of nematode genera, the data were submitted to the faunal analyses (Ferris et al., 2001) using the NINJA tool referred to as “an automated calculation system for nematode-based biological monitoring” (Sieriebriennikov et al., 2014). This way a graphical representation of the soil food web was obtained using enrichment and structural indices (EI and SI, respectively) (Ferris et al., 2001; Ferris, 2010).

### Soybean-maize cropping experiment

Student *t*-test (Statistica, Version 13.2; www.statsoft.com) analyses was done to determine whether significant ( $P \leq 0.05$ ) differences existed during both seasons between the two treatments (glyphosate-treated and non-treated plot halves) for the nematode population densities. Data were also subjected to one way analyses of variance (ANOVA) (Statistica Version 13.2)

to determine whether significant ( $P \leq 0.05$ ) differences existed for nematode population densities among the three sampling dates for both crops. In addition, terrestrial non-parasitic nematode data were subjected to faunal analyses using the program NINJA (Sieriebriennikov et al., 2014).

## Results

### Commercial soybean field study

Thirty-two non-parasitic nematode genera were collectively identified from soils of the three ecosystems, with 65% identified from soils of glyphosate-tolerant soybean fields, 72% from conventional soybean fields, and 88% from natural vegetation sites (Table 4). The genera identified were represented by different feeding groups and functional guilds, and included bacterivores, fungivores, predators, and omnivores.

The predominant non-parasitic nematodes from glyphosate-tolerant soybean sites for the 2011/12 season were *Aphelenchus*, *Acrobeles*, and *Acrobeloides* (Table 5). *Aphelenchus* occurred in soils from all of glyphosate-tolerant soybean sites while *Acrobeles* and *Acrobeloides* occurred in only 50%. For conventional soybean, the predominant genera were *Panagrolaimus*, *Acrobeloides*, and *Aphelenchus*. *Panagrolaimus* occurred at 75% of the sites, with *Acrobeloides* and *Aphelenchus* occurring in 50%. In soils from natural vegetation sites, the predominant genera were *Acrobeles*, *Aphelenchus*, and *Acrobeloides*. *Acrobeles* occurred at all sites, while *Aphelenchus* and *Acrobeloides* were found at 75% of the sites.

For the 2012/13 season the predominant genera identified from soils of glyphosate-tolerant soybean sites were *Aphelenchus*, *Acrobeles*, and *Eucephalobus* (Table 5). *Aphelenchus* occurred at all sites and *Acrobeles* and *Eucephalobus* at 89% and 78%, respectively. For conventional soybean, the predominant genera were *Aphelenchus*, *Eucephalobus*, and *Acrobeloides*. *Aphelenchus* occurred at all sites, with

**Table 5. Prominence values (PV), mean population density (MPD) and frequency of occurrence (FO%) of non-parasitic nematode genera identified from 200 g soil samples from glyphosate-tolerant and conventional soybean fields, as well as natural vegetation from 28 sites in the soybean production area of South Africa during the 2011/12 and 2012/13 growing seasons.**

Genus	PV	FO%	MPD	Genus	PV	FO %	MPD	Genus	PV	FO%	MPD
Glyphosate-tolerant soybean				Conventional soybean				Natural vegetation			
<b>2011/12 season</b>											
<i>Aphelenchus</i>	3,646	100	3,646	<i>Panagrolaimus</i>	2,723	75	3,144	<i>Acrobeles</i>	3,885	100	3,885
<i>Acrobeles</i>	2,131	50	3,014	<i>Acrobeloides</i>	1,963	50	2,804	<i>Aphelenchus</i>	3,741	75	4,320
<i>Acrobeloides</i>	982	50	1,403	<i>Aphelenchus</i>	1,683	50	2,380	<i>Acrobeloides</i>	2,758	75	3,185
<i>Eucephalobus</i>	668	25	1,335	<i>Aphelenchoides</i>	840	25	1,680	<i>Panagrolaimus</i>	1,310	50	1,853
<i>Aphelenchoides</i>	572	25	1,144	<i>Acrobeles</i>	444	50	634	<i>Eucephalobus</i>	907	50	1,281
<i>Psilenchus</i>	452	25	905	<i>Plectus</i>	338	25	675	<i>Rhabditis</i>	648	50	915
<i>Panagrolaimus</i>	253	25	505	<i>Cephalobus</i>	216	25	431	<i>Cephalobus</i>	570	25	1,139
<i>Rhabditis</i>	197	25	393	<i>Eucephalobus</i>	171	25	342	<i>Tylenchus</i>	534	75	615
<i>Mesorhabditis</i>	168	25	335	<i>Monhystera</i>	158	25	315	<i>Psilechus</i>	184	50	262
<i>Tylenchus</i>	17	25	34	<i>Mesorhabditis</i>	130	25	259	<i>Zeldia</i>	71	25	142
<i>Discolaimoides</i>	3	25	5	<i>Tylenchus</i>	40	25	80	<i>Dorylaimus</i>	5	25	9
<i>Thornenema</i>	3	25	5	<i>Psilenchus</i>	20	25	40	<i>Aporcellaimellus</i>	4	25	8
<i>Paraxonchium</i>	2	25	4	<i>Dorylaimus</i>	4	25	7	<i>Discolaimium</i>	3	25	5
–	–	–	–	<i>Discolaimoides</i>	3	25	5	<i>Discolaimoides</i>	2	25	3
–	–	–	–	<i>Paraxonchium</i>	2	25	3	–	–	–	–
–	–	–	–	<i>Thornenema</i>	2	25	3	–	–	–	–
<b>2012/13 season</b>											
<i>Aphelenchus</i>	2,275	100	2,275	<i>Aphelenchus</i>	1,448	100	1,448	<i>Aphelenchus</i>	2,590	100	2,590
<i>Acrobeles</i>	1,230	89	1,304	<i>Eucephalobus</i>	988	44	1,490	<i>Eucephalobus</i>	2,425	83	2,662
<i>Eucephalobus</i>	941	78	1,065	<i>Acrobeloides</i>	821	67	1,003	<i>Panagrolaimus</i>	1,615	100	1,615
<i>Acrobeloides</i>	827	67	1,010	<i>Panagrolaimus</i>	553	33	962	<i>Cephalobus</i>	1,414	67	1,728
<i>Rhabditis</i>	824	67	1,007	<i>Acrobeles</i>	365	33	635	<i>Acrobeloides</i>	1,156	83	1,269
<i>Aphelenchoides</i>	781	56	1,043	<i>Aphelenchoides</i>	334	22	711	<i>Acrobeles</i>	849	56	1,134
<i>Panagrolaimus</i>	576	56	770	<i>Seleborca</i>	286	22	610	<i>Aphelenchoides</i>	817	67	998
<i>Zeldia</i>	298	33	519	<i>Cephalobus</i>	587	56	784	<i>Ditylenchus</i>	323	33	562
<i>Cephalobus</i>	148	22	315	<i>Plectus</i>	241	22	513	<i>Plectus</i>	183	33	319
<i>Chiloplacus</i>	63	11	190	<i>Ditylenchus</i>	212	22	451	<i>Mesorhabditis</i>	151	17	365
<i>Ditylenchus</i>	53	11	161	<i>Mesorhabditis</i>	206	33	439	<i>Seleborca</i>	77	17	186
<i>Mesorhabditis</i>	42	11	126	<i>Plectus</i>	201	22	429	<i>Rhabditis</i>	68	17	165
<i>Plectus</i>	34	11	108	<i>Zeldia</i>	98	11	295	<i>Zeldia</i>	66	17	159
<i>Tylencholaimus</i>	7	11	22	<i>Chiloplacus</i>	78	11	235	<i>Wilsonema</i>	65	17	157
<i>Discolaimium</i>	3	22	7	<i>Dorylaimus</i>	8	55	11	<i>Chilopacus</i>	63	17	154
<i>Aporcellaimellus</i>	3	22	6	<i>Leptonchus</i>	4	11	11	<i>Teratocephalus</i>	7	17	16
<i>Eudorylaimus</i>	2	11	7	<i>Tylencholaimus</i>	3	11	10	<i>Eudorylaimus</i>	7	33	12

<i>Discolaimoides</i>	2	11	6	<i>Alaimus</i>	3	11	9	<i>Leptonchus</i>	5	17	13
–	–	–	–	<i>Mononchus</i>	2	11	5	<i>Coslenchus</i>	4	17	10
–	–	–	–	–	–	–	–	<i>Tylencholaimellus</i>	4	17	9
–	–	–	–	–	–	–	–	<i>Discolaimoides</i>	3	17	8

*Eucephalobus* and *Acrobeloides* occurring at 44% and 67%, respectively. In soils from natural vegetation sites the predominant genera were *Aphelenchus*, *Eucephalobus* and *Panagrolaimus*. *Aphelenchus* and *Panagrolaimus* occurred at all sites and *Eucephalobus* at 83%.

Although the abundance of the predominant genera (*Acrobeles*, *Acrobeloides*, *Aphelenchus*, *Eucephalobus*, and *Panagrolaimus*) varied substantially for the three ecosystems, it did not differ significantly between ecosystems according to *t*-Test analyses (Table 6).

Mixed Models analysis showed significant ( $P \leq 0.05$ ) interactions for fungivores, omnivores, and predators for Season\*Locality and for predators for Season\* Ecosystem\*Locality (Table 7). Due to relative low *F*-ratios for this interaction for fungivores, and the absence or very low numbers for predators and omnivores (ranging between 2 and 7 for omnivores and 2 and 4 for predators (Table 4) further discussion of the data is abstained from.

Season significantly ( $P \leq 0.05$ ) affected the abundance of all four nematode trophic groups (bacteri-, fungi-, omnivores, and predators) (Table 7). The abundance of bacterivores ( $873 \pm 426$  vs.  $120 \pm 430$  nematodes/200g soil), fungivores ( $283 \pm 150$  vs.  $88 \pm 152$  nematodes/200g soil) and omnivores ( $1.6 \pm 0.9$  vs.  $0.5 \pm 0.9$  nematodes/200g soil) was significantly higher in Season 2 compared with Season 1. By contrast, predator abundance was significantly ( $P \leq 0.05$ ) higher in Season 1 ( $0.8 \pm 0.15$  nematodes/200g soil) than Season 2 ( $0.38 \pm 0.12$  nematodes/200g soil). However, due to either the absence or very low numbers for predators and omnivores discussion of the data for these two trophic groups is abstained from.

Ecosystem affected only predator abundance significantly ( $P \leq 0.05$ ), with significantly higher population densities in glyphosate-tolerant ( $1 \pm 0.2$  nematodes/200g soil) compared with conventional soybean ( $0.3 \pm 0.2$  nematodes/200g soil) and natural veld ( $0.2 \pm 0.1$  nematodes/200g soil). However, the very low predator numbers recorded for all three ecosystems warrants no further discussion.

Locality significantly ( $P \leq 0.05$ ) affected omnivore and predator abundance but warrants no further discussion due to very low population densities recorded for these to trophic groups (Tables 5 and 7).

According to CCA analyses, no differences were apparent for the nematode assemblages present in soils from the three ecosystems when data for the sites were combined (data not shown). However, when the three ecosystems were plotted per site, distinct variations existed among the respective nematode communities for the three ecosystems with the cumulative explained variation (Axes 1 and 2) for the different locations for both seasons ranging from 22% to 82% (data not shown). An example is that of Edenville (Fig. 2) with a cumulative explained variation of 48.9%. For the other localities, similar differences between the nematode communities for the three ecosystems were observed (data not shown) although the nematode assemblages associated with each ecosystem differed among the localities.

According to faunal analysis, soils from the majority of the sites (54%) of the three ecosystems plotted in Quadrant D due to their Enrichment Index (EI) and Structural Index (SI) being <50% for both seasons (Fig. 3a). Such soils were dominated mainly by the presence of fungivores, especially Fu2. Forty-six percent of the sites plotted in Quadrant A due to their EI being >50% and SI being <50%. These soils were dominated by bacterivores, mainly belonging to Ba1 and Ba2. None of the sampling sites plotted in the Quadrants B and/or C.

The metabolic footprints (data pooled for sites from each ecosystem for each season), for the three ecosystems were small (Fig. 3b). The EI for the three ecosystems was intermediate (38%) to moderately high (68%) and the SI very low (<10%) for both seasons. Small differences were evident for both natural vegetation (plotted in Quadrant D for the two respective seasons) and glyphosate-tolerant (plotted in Quadrant A for the two respective seasons) ecosystems. However, for the conventional soybean ecosystem the difference for the two seasons was more pronounced, plotting in Quadrant A (2011/2012 growing season) and D (2012/2013 growing season). This phenomenon was probably due to a higher percentage of Fu2 being present in soils during the 2013 season.

### Soybean-maize cropping experiment

All nematode genera identified from the experimental plot were present in soil samples taken before the

**Table 6. Non-parasitic nematodes genera mean population density data per 200 g rhizosphere soil of glyphosate-tolerant and conventional soybean crops, as well natural vegetation from 28 sites surveyed in the soybean production areas of South Africa during the 2011/12 and 2012/2013 growing seasons. Values shown are means, followed by the standard deviation (SD).**

2011/12			2012/13		
Ecosystems	t-value	P	Ecosystems	t-value	P
<b><i>Acrobeles</i></b>					
Glyphosate-tolerant soybean: 603 ± 1,348	0.15	0.88	Glyphosate-tolerant soybean: 261 ± 583	0.06	0.97
Conventional soybean: 27 ± 284			Conventional soybean: 50 ± 335		
Glyphosate-tolerant: 603 ± 1,348	-0.02	0.98	Glyphosate-tolerant soybean: 261 ± 583	0.01	0.99
Natural vegetation: 777 ± 1,737			Natural veld: 227 ± 507		
Conventional soybean: 127 ± 284	-0.17	0.87	Conventional soybean: 150 ± 335	-0.04	0.97
Natural vegetation: 777 ± 1,737			Natural veld: 227 ± 507		
<b><i>Acrobelloides</i></b>					
Glyphosate-tolerant soybean: 281 ± 627	0.06	0.96	Glyphosate-tolerant soybean: 202 ± 452	0.001	1
Conventional soybean: 560 ± 1,254			Conventional soybean: 201 ± 449		
Glyphosate-tolerant: 281 ± 627	0.08	0.94	Glyphosate-tolerant: 202 ± 452	0.02	0.98
Natural vegetation: 637 ± 1,424			Natural vegetation: 254 ± 568		
Conventional soybean: 560 ± 1,254	-0.01	0.99	Conventional soybean: 201 ± 449	0.02	0.98
Natural vegetation: 637 ± 1,424			Natural vegetation: 254 ± 568		
<b><i>Aphelenchus</i></b>					
Glyphosate-tolerant soybean: 729 ± 1,631	-0.04	0.97	Glyphosate-tolerant soybean: 455 ± 1,017	0.04	0.97
Conventional soybean: 476 ± 1,064			Conventional soybean: 290 ± 648		
Glyphosate-tolerant: 729 ± 1,631	-0.12	0.91	Glyphosate-tolerant: 455 ± 1,017	0.01	0.99
Natural vegetation: 864 ± 1,932			Natural vegetation: 518 ± 1,158		
Conventional soybean: 476 ± 1,064	-0.05	0.96	Conventional soybean: 290 ± 648	0.05	0.96
Natural vegetation: 864 ± 1,932			Natural vegetation: 518 ± 1,158		
<b><i>Eucephalobus</i></b>					
Glyphosate-tolerant soybean: 267 ± 597	-0.15	0.89	Glyphosate-tolerant soybean: 213 ± 476	-0.03	0.97
Conventional soybean: 68 ± 153			Conventional soybean: 298 ± 666		
Glyphosate-tolerant: 267 ± 597	-0.004	1	Glyphosate-tolerant: 213 ± 476	0.09	0.93
Natural vegetation: 256 ± 573			Natural vegetation: 532 ± 1,191		
Conventional soybean: 68 ± 153	-0.14	0.89	Conventional soybean: 298 ± 666	0.05	0.96
Natural vegetation: 256 ± 573			Natural vegetation: 532 ± 1,191		
<b><i>Panagrolaimus</i></b>					
Glyphosate-tolerant soybean: 101 ± 226	0.18	0.86	Glyphosate-tolerant soybean: 154 ± 344	-0.02	0.98
Conventional soybean: 629 ± 1,406			Conventional soybean: 192 ± 430		
Glyphosate-tolerant: 101 ± 226	0.13	0.9	Glyphosate-tolerant: 154 ± 344	0.07	0.94
Natural vegetation: 371 ± 829			Natural vegetation: 323 ± 722		
Conventional soybean: 629 ± 1,406	0.05	0.96	Conventional soybean: 192 ± 430	0.05	0.96
Natural vegetation: 371 ± 829			Natural vegetation: 323 ± 722		

**Table 7. Significance values (P and F-ratios) for three independent variables (ecosystem, locality, and season), according to a Mixed Models analysis, showing their effects (individually and in combination) on four non-parasitic nematode trophic groups that were identified in the soybean production areas of South Africa during the 2011/12 and 2012/13 growing seasons.**

Source	Bacterivores		Fungivores		Omnivores		Predators	
	F	P	F	P	F	P	F	P
Season	42.158	0.001**	17.322	0.001**	10.078	0.006**	4.531	0.049**
Ecosystem	0.489	0.622	0.536	0.595	0.675	0.523	6.384	0.009**
Season*Ecosystem	0.123	0.885	0.021	0.980	0.300	0.745	1.238	0.316
Locality	1.315	0.307	2.494	0.075	4.851	0.007**	8.648	0.001**
Season*Locality	1.073	0.388	3.540	0.039**	8.641	0.001**	24.999	0.001**
Ecosystem*Locality	0.580	0.794	0.930	0.526	0.554	0.814	2.001	0.108
Season*Ecosystem*Locality	0.561	0.728	1.454	0.259	1.076	0.410	7.083	0.001**

\*Indicates interaction between and among independent variables; \*\*Denotes significance at  $P < 0.05$  according to the Mixed Modes analysis (SPSS, Version 25).




Figure 2: Canonical correspondence analyses (CCA) of population densities of terrestrial, non-parasitic nematode genera identified from 200g soil samples obtained from three ecosystems: glyphosate-tolerant soybean fields, conventional soybean fields, and natural vegetation in Edenville during 2012/13 season.


study commenced. Their numbers were, however, low and ranged between two and seven per 200g soil.

Fourteen non-parasitic nematode genera were identified from rhizosphere soil samples. In general, higher numbers of non-parasitic nematodes were recorded during the 2014/15 compared with the 2013/14 growing season (Table 8). *Aphelenchus* was most abundant and always occurred in higher numbers in glyphosate-treated plots. *Aphelenchoides* only occurred in the glyphosate-treated half of the plot while *Tylenchus* only occurred in non-treated halves of both crops. *Acrobeloides*, *Cephalobus*, *Eucephalobus*, and *Panagrolaimus* always occurred in higher numbers in the glyphosate-treated compared with the non-treated half of the plots.

### Faunal analysis

Substantial differences were apparent for non-parasitic nematode assemblages present in soils of the soybean-maize cropping system for the glyphosate-treated (plotted below the red line in Fig. 4) compared with the non-treated plot halves (plotted above the red line in Fig. 4). Data for the non-treated soil of all sampling dates plotted in Quadrants A and B, with EI >45% due to domination by bacterivores (Ba2 in particular representing *Acrobeles*, *Acrobeloides*, and *Eucephalobus*). One sample from the non-treated maize plants plotted in Quadrant B with a high SI (86%) due to the presence of predators (Pr5)

## Non-parasitic nematodes of grains



Figures 3a. & 3b. Faunal profiles (Sieriebriennikov *et al.*, 2014) representing the enrichment and structural conditions of soil food webs on the abundance and diversity of terrestrial, non-parasitic nematode genera identified from soils of glyphosate-tolerant and conventional soybean fields, as well as adjacent natural vegetation sites (39 in total) sampled during 2011/12 and 2012/13 seasons (A) and data for such sites pooled for the two seasons (B) in South African soybean production areas. The rhombus solid line around the mean indicates the metabolic footprint, the dotted line indicates the deviation of the metabolic footprint.

belonging to the genera *Aporcelaimellus* and *Discolaimium*. By contrast, all samples from the glyphosate-treated plot half, except for one, plotted in Quadrants C and D with a low EI (<35%). This was substantiated by the presence of fungivores, Fu2 in particular belonging to *Aphelenchus*, and *Aphelenchoides* while Fu4 was also present and represented by *Tylencholaimus*.

## Discussion

The 32 non-parasitic nematode genera identified from the commercial soybean field study and adjacent vegetation, and an experimental site where a soybean-maize rotation was done represent novel information for South Africa. Previous studies in such agricultural areas only focused on plant-parasitic nematodes (Rieker and Henshaw, 1998; Fourie *et al.*, 2001).

Various abiotic factors are known to impact on nematode development and survival (Perry *et al.*, 2013), with season significantly shown to affect the abundance of the four non-parasitic nematode trophic groups recorded in our study. This scenario implies that prevailing environmental conditions played a pronounced role during the two seasons this study was conducted.

Although soils from the commercial glyphosate-tolerant fields were dominated by the fungivore genus *Aphelenchus* during both seasons of the study, this genus also dominated in soils from conventional soybean and natural vegetation ecosystems in the second season. In the soybean-maize cropping experiment, it dominated in the second season in both plots. These results agree with those by Neher *et al.* (2014) who recorded higher abundance of fungivores in soils from

Bt maize compared with those from their near-isolines. Also, it is to a certain extent in agreement with those by Liphadzi *et al.* (2005) who stated that fungivores dominated in soils treated with various herbicides. These authors, however, did not refer to glyphosate-treated soils as was done in the present studies.

The abundance and dominance of the non-parasitic nematode genera, however, varied among the three ecosystems sampled during the extensive field study, and for the 2-year experimental soybean-maize cropping study. For the field study, the glyphosate-tolerant soybean ecosystems supported the least number of genera (21), while the natural vegetation supported the most (27), followed by the conventional soybean ecosystem (23). This trend is in agreement with reports by Bekker (2016) that natural vegetation ecosystems adjacent to maize fields in South Africa supported a higher diversity of non-parasitic nematodes than conventional and conservation maize ecosystems. Also, the general trend that nematode communities in soybean fields and natural vegetation sites were dominated by bacterivore genera of the families Acrobelidae, Cephalobidae, and Panagrolaimidae and fungivores of the families Aphelenchidae and Aphelenchoididae is in agreement with results by Bekker (2016) who did a similar study for commercial maize fields. The dominance of bacterivores in terms of the genera diversity in soils sampled during the present studies is also in agreement with reports by Djigal *et al.* (2004) and Xu *et al.* (2015). These authors suggested that bacterial feeding nematodes are the most abundant metazoans in soil substrates.

Fungivores were the second most prevalent group in soils sampled in the present studies, which is in

**Table 8. Number of non-parasitic nematodes per 200 g rhizosphere soil of soybean cv. LS 6164 R and maize cv. DKC 80–30 RR plants in glyphosate-treated and non-treated small-field plot halves at three sampling dates during the 2013/14 and 2014/15 growing seasons. Values shown are means, followed by the standard deviation (SD).**

Nematode genus	Functional guild, followed by c-p value	Soybean				Maize			
		Glyphosate-treated	Non-treated	t-value	P	Glyphosate-treated	Non-treated	t-value	P
<i>Acrobeles</i>	Ba2	–	298±527	–7	<b>0.001</b>	888±1,029	–	–8.15	<b>0.001</b>
<i>Acrobelloides</i>	Ba2	201±342	155±172	0.9	0.37	1,001±1,795	149±169	–1.57	0.12
<i>Aphelenchus</i>	Fu2	475±746	189±249	0.31	0.76	2,635±3,020	1,762±2,002	–6.68	0.5
<i>Aphelenchoides</i>	Fu2	240±394	–	5.25	<b>0.001</b>	1,226±2,090	–	–4.15	<b>0.001</b>
<i>Aporcelaimellus</i>	Pr5	–	0.17±1	–1	0.33	–	–	–	–
<i>Cephalobus</i>	Ba2	512±1,011	189±298	3.73	<b>0.001</b>	745±1,098	256±316	–0.96	0.34
<i>Discolaimium</i>	Pr5	–	0.22±1	–0.99	0.33	–	–	–	–
<i>Ditylenchus</i>	Fu2	38±92	–	3.16	<b>0.002</b>	–	–	–	–
<i>Eucephalobus</i>	Ba2	211±407	46±20	0.81	0.42	1,000±1,129	89±156	–3.98	<b>0.001</b>
<i>Leptonchus</i>	Fu4	10±17	5±13	1.54	0.13	–	–	–	–
<i>Panagrolaimus</i>	Ba2	544±1,002	–	17.65	<b>0.001</b>	1,350±1,542	267±467	–3.41	<b>0.001</b>
<i>Teratocephalus</i>	Ba4	–	–	–	–	2±7	16±16	5.46	<b>0.001</b>
<i>Tylenchus</i>	Fu2	–	306±537	–4.1	<b>0.001</b>	–	267±462	4.16	<b>0.001</b>
<i>Tylencholaimus</i>	Fu4	5±15	7±17	–0.02	0.99	24±35	12±23	–2.76	<b>0.007</b>

– No nematodes recovered. Functional guilds given according to Ferris et al. (2001); Colonizer-persister (c-p) values given according to Bongers (1990) with Ba, Bacterivores; Fu, Fungivores; and Pr, Predators.

agreement with a recent study by Renčo and Čerevková (2017). These authors reported that fungivores are the second most abundant in soil after bacterivores nematodes. The lower abundance and occurrence of predators and omnivores in the commercial field study was not surprising since these two groups are regarded as being very sensitive to soil disturbances (Ferris et al., 2001). A similar trend was reported by Bekker (2016) for a commercial maize field study. Hence, despite that ecosystem significantly affected predator abundance, the very low population densities and/or absence of this trophic group at various sites are suggested to have caused this effect and hence discussion of the data is abstained from. The absence of omnivores in soils of the 2-year experi-

mental soybean-maize study is another interesting observation and cannot be explained at this stage.

Nematode communities generally differ and fluctuate substantially among different locations in terms of abundance, diversity, and occurrence (Franco-Navarro & Godínez-Vidal, 2017). This tendency, although not significant, was apparent for the three ecosystems sampled during the commercial field study. When the three ecosystems were, however, analysed per site using the nematode trophic groups each generally had different nematode communities and was separated from each other according to CCA analyses. However, no trend existed where a specific nematode genus/genera was exclusively associated with either of the three ecosystems. Although it was not possible to deduct the impact




Figure 4: A faunal, soil food web profile representing the enrichment and structural indices (EI and SI, respectively) of terrestrial, non-parasitic nematode assemblages identified at the three sampling dates in glyphosate-tolerant and non-treated plot halves planted with soybean (during 2013/14 growing season) and maize (during 2014/15 growing season).

of each ecosystem on the nematode communities, our study showed that glyphosate-tolerant soybean had no deleterious effects on non-target beneficial nematodes. This is in agreement with those for other genetically-modified crops, for example, Al-Deeb et al. (2013) and Neher et al. (2014) demonstrating that genetically-modified, Bt maize had no significant adverse effects on non-target, beneficial, and plant-parasitic nematodes. Also, Chen et al. (2017) concluded that Bt rice had no remarkable impact on beneficial soil nematode communities and was pest specific. However, Neher et al. (2014) suggested that rhizosphere soil from Bt maize may contain more complex and successfully mature nematode communities opposed to those from non-Bt near isolines which may be applicable to our study also where fungivores generally dominated in soil from glyphosate-treated soybean crops. This phenomenon may be an indication that less disturbance in the glyphosate-treated soybean fields probably can contribute to nematode communities being more matured.

According to faunal analysis, all soybean sites sampled were disturbed and degraded, indicating that the quality of these soils is not optimal in terms of the presence of beneficial nematodes (Ferris et al., 2001). This situation is often associated with management practices such as repeated tillage (Berkelmans et al., 2003) and pesticide application (Carrascosa et al., 2014) which are typical practices in local soybean production areas (Liebenberg, 2012). Contrary to

annual crop fields, natural vegetation ecosystems are usually regarded as stable and structured due to either no or minimal disturbances (Ferris et al., 2001). However, in the current study all natural vegetation sites were also degraded or disturbed. This might be explained by the vegetation type that was represented by mainly grasses. Often natural vegetation consists of woody, perennial plants that are mostly considered less disturbed than grassland vegetation (Cullman et al., 2010). The latter vegetation probably experiences periods during which the organic content of the soil is high compared to periods when substantially less organic material is present (Shaw et al., 2016).

Results from the soybean-maize cropping experiment, however, showed that glyphosate applied as a leaf spray twice per season during two consecutive growing seasons generally affected the abundance and diversity of non-parasitic nematodes. This was substantiated by soil food web analysis of the different nematode sampling dates that showed that the majority of the glyphosate-treated plots for both seasons were degraded and depleted opposed to the non-treated plots that were disturbed but enriched. These results are not in agreement with those of the commercial field study and also those reported by Liphadzi et al. (2005), who found that glyphosate application had no effect on the abundance and diversity of non-parasitic nematodes in glyphosate-treated plots during a 3-year study.



It is worth mentioning that the non-treated plot, was hoed, implying some disturbance in the upper soil while organic material was also added to the soil. Both these activities might have had an effect on non-parasitic nematode communities and probably favoring bacterivore genera.

Ultimately, results from the two South African studies conducted showed similarity in terms of *Aphelenchus* domination. However, glyphosate application did not affect the general abundance of non-parasitic nematodes compared with those from conventional soybean fields and natural vegetation sites where no glyphosate had been applied for at least 5 years prior to this study or never before.

## Acknowledgments

This work is based on the research supported by the National Research Foundation under Grant SUR20110707000020249. Any opinion, finding, and conclusion or recommendation expressed in this material is that of the author(s) and the NRF does not accept any liability in this regard. Personnel and students of the NWU and ARC-GCI, in particular Dr. Suria Ellis (Biometrician, NWU), as well as all the farmers on which farms samples were taken are thanked for their technical assistance. The Agricultural Research Council, Institute for Soil, Climate and Water, AgroClimatology, is acknowledged for supplying weather data.

## References

- Al-Deeb, M., Wilde, G.E., Blair, J.M., and Todd, T.C. 2013. Effect of *Bt* corn for corn rootworm control on nontarget soil microarthropods and nematodes. *Environmental Entomology* 32: 859–65.
- Allegrini, M., Zabaloy, M.C., and Gomez, V. 2015. Ecotoxicological assessment of soil microbial community tolerance to glyphosate. *Science of the Total Environment* 533: 60–8.
- Bekker, S. 2016. Nematode communities: bio-indicators of soil quality in conventional and conservation agricultural cropping systems. PhD Dissertation, North-West University. Potchefstroom, South Africa.
- Beretta, A.N., Silbermann, A.V., Paladino, L., Torres, D., Bassahun, D., Musselli, R., and García-Lamohte, A. 2014. Soil texture analyses using a hydrometer: Modification of the Bouyoucos method. *Ciencia e Investigacion Agraria* 41: 263–71.
- Berkelmans, R., Ferris, H., Tenuta, M., and Van Bruggen, A.H.C. 2003. Effects of long-term crop management on nematode trophic levels other than plant feeders disappear after 1 year of disruptive soil management. *Applied Soil Ecology* 23: 223–35.
- Bongers, T. 1990. The maturity index: an ecological measure of environmental disturbance based on nematode species composition. *Oecologia* 83: 14–9.
- Bongers, T., and Bongers, M. 1998. Functional diversity of nematodes. *Applied Soil Ecology* 10: 239–51.
- Bouyoucos, G.J. 1962. Hydrometer method improved for making particle size analysis of soils. *Agronomy Journal* 54: 464–65.
- Carrascosa, M., Sánchez-Moreno, S., and Alonso-Prados, J.L. 2014. Relationships between nematode diversity, plant biomass, nutrient cycling and soil suppressiveness in fumigated soils. *European Journal of Soil Biology* 62: 49–59.
- Cerdeira, A.L., Gazziero, D.L.P., Duke, S.O., Matallo, M.B., and Spadotto, C.A. 2007. Review of potential environmental impacts of transgenic glyphosate-resistant soybean in Brazil. *Journal of Environmental Science and Health Part B* 42: 539–49.
- Cerdeira, A.L., and Duke, S.O. 2010. Effects of glyphosate-resistant crop cultivation on soil water quality. *GM Crops* 1: 16–24.
- Chen, X., Liu, T., Li, X., Li, H., Chen, F., Liu, M., and Whalen, T.C. 2017. Soil nematode community varies between rice cultivars but is not affected by transgenic *Bt* rice expressing *Cry1Ab* or *Cry1Ab/Cry1Ac*. *Biology and Fertility of Soils* 53: 501–9.
- Cullman, S.W., DuPont, S.T., Glover, J.D., Buckley, D.H., Fick, G.W., Ferris, H., and Crews, T.E. 2010. Long-term impacts of high-input annual cropping and unfertilized perennial grass production on soil properties and belowground food webs in Kansas, USA. *Agriculture Ecosystems and Environment* 137: 13–24.
- De Waele, D., and Jordaan, E.M. 1988. Plant-parasitic nematodes on field crops in South Africa. 1. *Maize. Revue de Nematologie* 11: 203–12.
- Dill, G.M., Cajacob, C.A., and Padgett, S.R. 2008. Glyphosate-resistant crops: adoption, use and future considerations. *Pest Management Science* 64: 326–31.
- Dijgal, D., Brauma, A., Diop, T. A., Chotte, J. L., and Villenave, C. 2004. Influence of bacterial-feeding nematodes (Cephalopidae) on soil microbial communities during maize growth. *Soil Biology and Biochemistry* 36: 323–31.
- Dlamini, T.S., Tshabalala, P., and Mutengwa, T. 2014. Soybeans production in South Africa. *Oilseeds and Fats Crops and Lipids* 21: D207.
- Doncaster, C.C., Edwards, B.S., and Shepherd, A.M. 1967. A labour-saving method of making Fenwick multi chamber counting slides for nematodes. *Nematologica* 12: 644.
- Duke, S.O., and Powles, S.B. 2008. Mini-review. Glyphosate: A once-in-a-century herbicide. *Pest Management Science* 64: 319–25.
- Ferris, H., Vanette, R. C., Van der Meulen, H. R. and Lau, S. S. 1998. Nitrogen mineralization by bacterial-feeding nematodes: Verification and measurement. *Plant and Soil* 203: 159–71.

- Ferris, H., Bongers, T., and De Goede, R.G.M. 2001. A framework for soil food web diagnosis: extension of the nematode faunal analysis concept. *Applied Soil Ecology* 18: 13–29.
- Ferris, H. 2010. Contribution of nematodes to the structure and function of the soil food web. *Journal of Nematology* 42: 63–7.
- Fourie, H., Mc Donald, A.H., and Loots, G.C. 2001. Plant-parasitic nematodes in field crops in South Africa. 6. Soybean. *Nematology* 3: 447–54.
- Franco-Navarro, F., and Godinez-Vidal, D. 2017. Soil nematodes associated with different land uses in the Los Tuxtlas Biosphere Reserve, Veracruz, Mexico. *Revista Mexicana de Biodiversidad* 88: 136–45.
- Fraschetti, S., Guarnieri, G., Bevilacqua, S., Terlizzi, A., and Danovaro, R. 2016. Impact of offshore gas platforms on the structural and biodiversity of nematodes. *Marine Environmental Research* 111: 56–64.
- Hooper, D.J., Hallman, J., and Subbotin, S. 2005. Methods for extraction, processing and detection of plant and soil nematodes, in Luc, M., Sikora, R.A., and Bridge, J. (Eds), *Plant parasitic nematodes in subtropical and tropical agriculture*, CAB International, United Kingdom, pp. 53–86.
- Hurley, T.M., Mitchell, P.D., and Frisvold, G.B. 2009. Weed management costs, weed best management practices, and the Roundup Ready® weed management program. *AgBioforum* 12: 281–290.
- James C. 2015. Global status of commercialized biotech/GM crops: 2015. ISAAA brief No. 51. Available at <http://www.isaaa.org/resources> (Accessed: 11/07/2017).
- Liebenberg, A. J. 2012. Sojaboon Produksiehandel-leiding/Soybean Production Manual. Agricultural Research Council-Grain Crops Institute. Potchefstroom, South Africa p. 159.
- Liphadzi, K.B., Al-Khatib, K., Bensch, C.N., Stahlman, P.W., Dille, J.A., Todd, T., Rice, C.W., Horak, M.J., and Head, G. 2005. Soil microbial and nematode communities as affected by glyphosate and tillage practices in a glyphosate-resistant cropping system. *Weed Science* 53: 536–45.
- Marais, M., Swart, A., Fourie, H., Berry, S.D., Knoetze, R., and Malan, A. 2017. Techniques and Procedures, in Fourie, H., Spaull, V.W., Jones, R.K., Daneel, M.S., and De Waele, D. (Eds), *Nematology in South Africa: A view from the 21st Century*, Springer Publishing, Germany, pp. 73–118.
- Neher, D.A. 2001. Role of nematodes in soil health and their use as indicators. *Journal of Nematology* 33: 161–68.
- Neher, D.A., Muthumbi, A.W.N., and Dively, G.P. 2014. Impact of coleopteran-active *Bt* on non-target nematode communities in soil and decomposing corn roots. *Soil Biology and Biochemistry* 76: 127–35.
- Newman, M.M., Hoilett, N., Lorenz, N., Dick, R.P., Liles, M.R., Ramsier, C., and Kloepper, J.W. 2016. Glyphosate effects on soil rhizosphere-associated bacterial communities. *Science of The Total Environment* 543: 155–60.
- Noel, G.R., and Wax, L.M. 2009. *Heterodera glycines* population development on soybean treated with glyphosate. *Nematropica* 39: 247–53.
- Osman, A.A., and Viglierchio, D.R. 1981. Herbicide effects in nematode diseases. *Journal of Nematology* 13: 417–19.
- Perry, R.N., Wright, D.J., and Chitwood, D.J. 2013. Reproduction, physiology and biochemistry, in Perry, R.N., and Moens, M. (Eds), *Plant Nematology*, 2nd ed., CAB International, United Kingdom, pp. 219–45.
- Renčo, M. and Čerevková, A. 2017. Windstorms as mediator of soil nematode community changes: Evidence from European spruce forest. *Helminthologia* 54: 36–47.
- Riekert, H.F., and Henshaw, G.E. 1998. Effect of soybean, cowpea and groundnut rotation on root-knot nematode build-up and infestation of dryland maize. *African Crop Science Journal* 6: 377–83.
- Shaw, E.A., Denef, K., De Tomasel, C.M., Cofruto, M.F., and Wall, D.H. 2016. Fire affects root decomposition, soil food web structure, and carbon flow in tall-grass prairie. *Soil* 2: 199–210.
- Shütte, G., Eckertorfer, M., Rastelli, V., Reichenbecher, W., Restrepo-Vassalli, S., Ruohonen-Lehto, W., Saucy, A., G., W., and Mertens, M. 2017. Herbicide resistance and biodiversity: Agronomic and environmental aspects of genetically modified herbicide-resistant plants. *Environmental Sciences Europe* 29: 5, <https://doi.org/10.1186/s12302-016-0100-y>.
- Sieriebriennikov, B., Ferris, H., and De Goede, R.G.M. 2014. NINJA: An automated calculation system for nematode-based biological monitoring. *European Journal of Soil Biology* 61: 90–3.
- Vega, J.J., Talatala-Sanico, R.L., and Gapasin, R.M. 1993. Effect of herbicides on the pathogenicity of root-knot nematode (*Meloidogyne incognita*) on soybean *Glycine max* (L.) Merr and *Amaranthus spinosus* L. Phillipine. *Phytopathology* 29: 42–53.
- Wada, S., Toyota, K., and Takada, A. 2011. Effects of nematicide imicyafos on soil nematode community structure and damage of radish caused by *Pratylenchus penetrans*. *Journal of Nematology* 43: 1–6.
- Walkey, A., and Black, I.A. 1947. A critical examination of a rapid method for determining organic carbon in soils-effect of variations in digestion conditions and of inorganic soil constituents. *Soil Science* 63: 251–64.
- Wardle, D.A., Williamson, W.M., Yeates, G.W., and Bonner, K.I. 2005. Trickle-down effects of aboveground trophic cascades on the soil food web. *Oikos* 111: 348–58.
- Xu, L., Wensi Xu, W., Ying Jiang, Y., Hu, F., and Li, H. 2015. Effects of interactions of auxin-producing bacteria and bacterial-feeding nematodes on regulation of peanut growths. *PLoS ONE* 10:e0124361.
- Yang, X., Harrison, S.K., and Riedel, R.M. 2002. Soybean (*Glycine max*) response to glyphosate and soybean cyst nematode (*Heterodera glycines*). *Weed Technology* 16: 332–39.
- Zhao, J., Neher, D.A., Fu, S., and Li, Z. 2013. Non-target effects of herbicides on soil nematode assemblages. *Pest Management Science* 69: 679–84.